

# A REMARK ON TRANSITIVITY OF GALOIS ACTION ON THE SET OF UNIQUELY DIVISIBLE ABELIAN EXTENSIONS IN $\text{Ext}^1(E(\overline{\mathbb{Q}}), \Lambda)$

MISHA GAVRILOVICH  
BALLIOL COLLEGE, OXFORD

ABSTRACT. We study Galois action on  $\text{Ext}^1(E(\overline{\mathbb{Q}}), \mathbb{Z}^2)$  and interpret our results as partially showing that the notion of a path on a complex elliptic curve  $E$  can be characterised algebraically. The proofs show that our results are just concise reformulations of Kummer theory for  $E$  as well as the description of Galois action on the Tate module.

Namely, we prove (a),(b) below by showing they are equivalent to (c) which is well-known: (a) group  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  acts transitively on the set of uniquely divisible abelian  $\text{End} E$ -module extensions of  $E(\overline{\mathbb{Q}})$  of algebraic points of an elliptic curve, by  $\Lambda \cong \mathbb{Z}^2$ , (b) natural algebraic properties characterise uniquely the Poincaré's fundamental groupoid of a complex elliptic curve, restricted to the algebraic points, (c) up to finite index, the image of the Galois action on the sequences  $(P_i)_{i>0}, jP_{ij} = P_i, i, j > 0$  of points  $P_i \in E^k(\overline{\mathbb{Q}})$  is as large as possible with respect to linear relations between the coordinates of the points  $P_i$ 's.

Our original motivations come from model theory.

...et on ajoutera: «je ne sais pas le reste»

## 1. INTRODUCTION

**1.1. Abelian group extensions.** The universal covering space of an elliptic curve  $A = E$  is just  $\mathbb{C}$ ; the linear structure on  $\mathbb{C}$  plays an important rôle in the theory of (complex) elliptic curves as algebraic varieties. This suggests that the relevant linear structure on  $\mathbb{C}$  is determined up to isomorphism by the algebraic curve itself. A uniquely divisible abelian extension of the group  $E(\mathbb{C})$  with kernel  $\mathbb{Z}^2$  describes such a linear structure; thus, the above considerations suggest that such an extension is unique, up to an automorphism  $\text{Aut}(\mathbb{C}/\mathbb{Q})$ .

The next proposition partially confirms these suggestions. We conjecture it holds for  $\mathbb{C}$  and in general, any algebraically closed field of zero characteristic.

**Proposition 1.** *Let  $E$  be an elliptic curve defined over a number field  $k \subset \overline{\mathbb{Q}}$ . Assume that all the endomorphisms of  $E$  are definable over  $k$ . If  $\text{End}E = \mathbb{Z}$ , then the Galois group  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  acts transitively on the set of uniquely divisible abelian extensions  $\text{Ext}_{\text{AbGroups}}^1(E(\overline{\mathbb{Q}}), \Lambda)$  of  $E(\overline{\mathbb{Q}})$  by  $\Lambda \cong \mathbb{Z}^2$ . If  $\text{End}E \neq \mathbb{Z}$ , then there are at most finitely many orbits on the set of uniquely divisible  $\text{End}E$ -module extensions in  $\text{Ext}_{\text{End}E\text{-mod}}^1(E(\overline{\mathbb{Q}}), \Lambda)$  of  $\text{End}E$ -modules.*

Here  $\Lambda$  denotes the kernel of  $\text{End}E$ -module covering map  $\mathbb{C} \rightarrow E(\mathbb{C})$ . We prove the proposition by an inductive argument using Kummer theory of  $E$  and the description of the Galois action on Tate module; moreover, the proof shows that Proposition 1 is equivalent to these arithmetic results.

**1.2. An algebraic notion of a path up to homotopy.** We ask whether the notion of paths (up to fixed point homotopy) on an elliptic curve  $E(\mathbb{C})$  may be described by its natural algebraic properties. “Paths up to fixed point homotopy” are usually thought of in the context of the Poincaré’s fundamental groupoid, which can be thought of as a 2-functor. Hence, we may reformulate the question as: *Is the fundamental groupoid functor on the complex algebraic varieties determined by its natural algebraic properties up to natural equivalence and an automorphism of the source category?* Proposition 2 is a partial positive answer to this question.

**Proposition 2** (Universality of fundamental groupoid functor). *Let  $E$  and a number field  $k$  be as above. Let  $\mathfrak{E}$  be the full subcategory of  $\text{Var}/\overline{\mathbb{Q}}$  consisting of Cartesian powers of  $E$ .*

*Let  $\underline{\Omega} : \mathfrak{E} \rightarrow \mathbf{Groupoids}$  be a functor satisfying conditions (1)-(3) of Definition 1 below (unique path-lifting along étale morphisms, preservation of direct product, etc).*

*Assume further that*

- (4)  $\underline{\Omega}(E)$  is a connected groupoid
- (5) there is an isomorphism

$$\Omega_{0,0}(E) := \{\gamma \in \Omega(E) : s(\gamma) = t(\gamma) = 0\} \cong \mathbb{Z}^2 \text{ as } \text{End}E\text{-modules.}$$

*Then, for  $E$  with no complex multiplication, there exists an automorphism  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that the functors  $\underline{\Omega}, \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma : \mathfrak{E} \rightarrow \mathbf{Groupoids}$  are naturally equivalent:*

$$\underline{\Omega} \cong \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma$$

*If  $E$  has complex multiplication, then there exists finitely many functors  $\mathbb{F}_1, \dots, \mathbb{F}_n : \mathfrak{E} \rightarrow \mathbf{Groupoids}$  satisfying conditions (1)-(5) above such*

that for any functor  $\underline{\Omega}$  satisfying (1)-(5) above there exists a Galois automorphism  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  and a number  $0 < i < n + 1$  such that  $\underline{\Omega}$  and  $\mathbb{F}_i \circ \sigma$  are naturally equivalent.

Here  $\text{Pt}_{\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}}(V) = V(\overline{\mathbb{Q}})$  and

$$\Omega_{\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}}(V) = \{\gamma \in \pi_1^{\text{top}}(V(\mathbb{C})) : s(\gamma), t(\gamma) \in V(\overline{\mathbb{Q}})\}$$

is the restriction of  $\pi_1^{\text{top}}$  to  $\overline{\mathbb{Q}}$ -rational points.

We prove Proposition 2 by reducing it to Proposition 1.

**1.3. Kummer theory and Galois representations.** We prove Propositions 1 and 2 by a rather simple induction based on Kummer theory and the image of Galois action on Tate module. The full proof is carried out in §3; here we just present an argument showing how these arithmetic results may be relevant. For simplicity assume  $\text{End} E = \mathbb{Z}$  and consider an extension  $H \rightarrow^\varphi E(\overline{\mathbb{Q}})$ . Take a pair  $\lambda_1, \lambda_2 \in \Lambda$  of generators of abelian group  $\Lambda = \ker \varphi$ ; the extension then gives rise to the infinite sequences  $(\varphi(\lambda_1/n))_n, (\varphi(\lambda_2/n))_n$  which can be thought of as elements of Tate module  $\prod_i T_i(E)$ . If we take another extension  $H' \rightarrow^{\varphi'} E(\overline{\mathbb{Q}})$ , we can obtain similar sequences  $((\varphi'(\lambda'_1/n))_n, (\varphi'(\lambda'_2/n))_n$ . If the extensions  $H \rightarrow^\varphi E(\overline{\mathbb{Q}})$  and  $H' \rightarrow^{\varphi' \circ \sigma} E(\overline{\mathbb{Q}})$  are isomorphic for some  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  and an isomorphism  $' : H \rightarrow H'$  takes  $(\lambda_1, \lambda_2)$  to  $(\lambda'_1, \lambda'_2)$ , then these pairs of sequences are conjugated by Galois action. Were  $\text{Gal}(\overline{\mathbb{Q}}/k) = \text{SL}_2(\hat{\mathbb{Z}})$ , it would be so for any two linearly independent pairs  $(\lambda_1, \lambda_2)$  and  $(\lambda'_1, \lambda'_2)$  of generators.

If  $\text{Gal}(\overline{\mathbb{Q}}/k)$  is of finite index in  $\text{SL}_2(\hat{\mathbb{Z}})$ , then it is sufficient to take  $\lambda'_1, \lambda'_2$  such that  $\varphi'(\lambda'_1/N) = \varphi(\lambda_1/N), \varphi'(\lambda'_2/N) = \varphi(\lambda_2/N)$ , for some  $N$  large enough; that is possible due to surjectivity of  $\varphi'$  and the unique divisibility of  $H'$ .

**1.4. Model theory: motivations and generalisations.** The results of this note appear naturally in a model-theoretic framework of «logically perfect structures» developed by Zilber [Zilb]; in this way they have been obtained in the author's DPhil thesis [Gav05]. We believe that the context and techniques of model theory are essential to generalise our results to other varieties and fields of arbitrary cardinality including the field of complex numbers. However, in this note we do not discuss the model theoretic motivations and techniques; we refer to the the author's DPhil thesis [Gav05] for that. The note presents only those results of the thesis which can be stated and proven in a model-theory free fashion; in particular, *the note omits the discussion of the Shafarevich conjecture on holomorphic convexity of universal covering*

*spaces and the case semi-Abelian varieties of higher dimension.* The results follow a line originated by Zilber [Zila] treating fully the case of the multiplicative group of an algebraically closed field of characteristic 0.

We would appreciate references to any literature discussing similar questions.

**1.5. Structure of the paper.** We state our results in detail in §2.

We prove Proposition 1 in §3. We establish the equivalence of the algebraic approach of Proposition 1 and the topological approach of Proposition 2 in §4. We state a precise conjecture about Shimura curves in §5.

## 2. RESULTS

In this section we state our results in full, and hint on a connection between the reformulations.

**2.1. Uniquely divisible extensions of Abelian groups.** Let  $E$  be an elliptic curve defined over a number field  $k \subset \overline{\mathbb{Q}} \subset \mathbb{C}$ , and let

$$0 \longrightarrow \Lambda \longrightarrow \mathbb{C} \xrightarrow{p} E(\mathbb{C}) \longrightarrow 0.$$

be the universal covering of  $E(\mathbb{C})$ . Let  $0 \in E(k)$  denote a  $k$ -rational point which is zero of the additive group  $E(\mathbb{C})$ .

**Proposition 1.** *Assume that all the endomorphisms of  $E$  are definable over  $k$ . If  $\text{End} E = \mathbb{Z}$ , then the Galois group  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  acts transitively on the set of uniquely divisible abelian extensions  $\text{Ext}_{\text{Ab Groups}}^1(E(\overline{\mathbb{Q}}), \Lambda)$  of  $E(\overline{\mathbb{Q}})$  by  $\Lambda \cong \mathbb{Z}^2$ . If  $\text{End} E \neq \mathbb{Z}$ , then there are at most finitely many orbits in uniquely divisible extensions in  $\text{Ext}_{\text{End} E\text{-mod}}^1(E(\overline{\mathbb{Q}}), \Lambda)$  of  $\text{End} E$ -modules.*

We conjecture the proposition holds for any algebraically closed field of zero characteristic, i.e. the action of  $\text{Aut}(\bar{K}/\mathbb{Q})$  on the set of uniquely divisible abelian extensions in  $\text{Ext}^1(E(\bar{K}), \Lambda)$ . Zilber [Zila] proves the transitivity of the  $\text{Aut}(\bar{K}/\mathbb{Q})$ -action on  $\text{Ext}^1(\bar{K}^*, \mathbb{Z})$  for arbitrary algebraically closed field  $K = \bar{K}$  of characteristic 0.

Remark that the set of non-equivalent extensions  $\text{Ext}_{\text{End} E\text{-mod}}^1(E(\overline{\mathbb{Q}}), \Lambda)$  is of cardinality  $2^{\aleph_0}$ . Also note that the injectivity of the profinite completion  $\hat{\Lambda}$  of the kernel  $\Lambda$  implies  $\text{Ext}_{\widehat{\text{End} E\text{-mod}}}^1(E(\overline{\mathbb{Q}}), \hat{\Lambda}) = 0$  making the proposition trivially true in this case.

A way to think of it is that the proposition claims that *it is possible to describe the universal covering space of an elliptic curve in a purely algebraic way*, admittedly with respect to a rather weak, linear structure on it.

**2.2. Relation between Propositions 1 and 2.** Here we interpret Proposition 1 as a universality property of the Poincaré fundamental groupoid functor. Identify points in  $\mathbb{C}$  and homotopy classes of paths in  $E(\mathbb{C})$  starting at 1 via the period map  $\gamma \mapsto \int_{\gamma} dz$ . Then the addition on  $\mathbb{C}$  corresponds to point-wise addition of paths, and dividing by  $n$  corresponds to path-lifting along the étale morphism  $nx : E(\mathbb{C}) \rightarrow E(\mathbb{C})$ . Using this correspondence, we interpret the unique divisibility of  $\mathbb{C}$  as the unique path-lifting property along étale morphisms from  $E(\mathbb{C})$  to  $E(\mathbb{C})$ .

These observations allow us to reformulate the previous result as an algebraic characterisation of the Poincaré's fundamental groupoid of a complex elliptic curve, i.e. that *the notion of paths (up to fixed point homotopy) on a complex elliptic curve  $E(\mathbb{C})$  may be described by its natural algebraic properties.*

A precise reformulation of the question is: *Is the fundamental groupoid functor on the complex algebraic varieties determined by its natural algebraic properties up to natural equivalence and an automorphism of the source category?* Proposition 2 provides a partial positive answer to this question; the proposition is just an equivalent reformulation of Proposition 1.

**2.3. The universality property of the Poincaré's fundamental groupoid functor.** In this subsection we introduce in detail the notions of category theory appropriate to state Proposition 2.

**2.3.1. The example: fundamental groupoid functor  $\pi_1^{\text{top}}(E(\mathbb{C}))$  as a two-functor.** It is convenient to consider 2-functors instead of functors to groupoids; the notions are equivalent. Before defining a 2-functor formally, we illustrate the notion by an example of Grothendieck [Gro].

The path 2-functor  $\underline{\Omega}$  on the category  $\mathfrak{Top}$  of topological spaces is a tuple  $(\text{Pt}, \Omega, s, t, \cdot)$  consisting of a *functor of points*  $\text{Pt} : \mathfrak{Top} \rightarrow \mathfrak{Sets}$  and a *paths functor*  $\Omega : \mathfrak{Top} \rightarrow \mathfrak{Sets}$  together with the following data:

- (1)  $\text{Pt}(T)$  is the set of points of topological space  $T$  and the morphism  $\text{Pt}(f) : \text{Pt}(T_1) \rightarrow \text{Pt}(T_2)$  is  $f : T_1 \rightarrow T_2$  as a map of sets.
- (2)  $\Omega(T)$  is the set of all paths in topological space  $T$ , i.e. continuous functions  $\gamma : [a, b] \rightarrow T$ ,  $a, b \in \mathbb{R}$ ; similarly  $\Omega(f), f \in \text{Hom}_{\mathfrak{Top}}(T_1, T_2)$  is the map taking a path  $\gamma : [a, b] \rightarrow T_1$  into  $f \circ \gamma : [a, b] \rightarrow T_2$ .
- (3)  $s_T, t_T : \Omega(T) \rightarrow \text{Pt}(T)$  are functions from the set of paths in  $T$  to their endpoints in  $T$ ; the function  $s(\gamma) = \gamma(a)$  (*source*) gives the beginning point of a path  $\gamma$ , and the function  $t(\gamma) = \gamma(b)$  (*target*) gives the ending point of path  $\gamma$ .

- (4)  $\cdot_T : \Omega(T) \times \Omega(T) \rightarrow \Omega(T)$  is the partial operation of the concatenations of paths, taking  $\gamma_1 : [a, b] \rightarrow T$ ,  $\gamma_2 : [b, c] \rightarrow T$  into  $\gamma = \gamma_1 \gamma_2$ ,  $\gamma|_{[a,b]} = \gamma_1$ ,  $\gamma|_{[b,c]} = \gamma_2$ .

Thus, a 2-functor from  $\mathfrak{Top}$  to  $\mathfrak{Sets}$  consists of two functors  $\text{Pt}, \Omega : \mathfrak{Top} \rightarrow \mathfrak{Sets}$ , and two natural transformations  $s, t : \Omega \rightarrow \text{Pt}$  from functor  $\Omega$  to  $\text{Pt}$ ;  $s$  stands for source and  $t$  stands for target. For each  $T$ , there is also a functorial associative operation  $\cdot_T$  defined on  $\Omega_{x,y}(T) \times \Omega_{y,z}(T) \rightarrow \Omega_{x,z}(T)$ , where  $\Omega_{x,y}(T) = \{\gamma \in \Omega(T) : s(\gamma) = x, t(\gamma) = y\}$ , etc; the operation  $\cdot$  makes  $\Omega_{x,x}(T)$  into a group; in the example above,  $\Omega_{x,x}(T)$  is the set of all loops in  $T$  based at  $x \in T$ .

In particular, for each  $T$  the set  $\Omega(T)$  carries the structure of a groupoid; in fact, it is conventional to consider  $\underline{\Omega}$  as a functor to the category of groupoids.

If in item (2) we define  $\Omega(T)$  to be the set of all paths *up to fixed point homotopy*, then we obtain the notion of the fundamental groupoid functor. The advantage of the original definition is that one may try and define  $n$ -functors describing  $n$ -dimensional homotopies on topological space  $T$ , cf. Grothendieck [Gro] for motivations; Voevodsky-Kapranov [VK91] propose an exact definition. Grothendieck [Gro] explains that it is essential not to insist on strict associativity etc, but rather to consider *all the identities to hold up to a homotopy of higher dimension*. This may be useful to generalise Proposition 2.

**2.3.2. Abstract fundamental groupoid functors.** Here we define the path-lifting property for a 2-functor, and an abstract fundamental groupoid functor as a functor preserving direct products, possessing the path-lifting property and with a particular functor of points.

**Definition 1.** *Let  $\underline{\Omega}$  be a 2-functor from a subcategory  $\mathfrak{E}$  of the category of varieties over an algebraically closed field  $K$ , into  $\mathfrak{Sets}$ . We say  $\underline{\Omega}$  is an abstract fundamental groupoid  $K$ -valued functor if  $\underline{\Omega}$  satisfies the following properties:*

- (1) *the functor  $\text{Pt}$  is the functor of  $K$ -rational points:*

$$\text{Pt}(X) = X(K) = \mathcal{M}or_{\text{Var}/K}(0, X)$$

*(here  $0$  denotes a single point variety defined over  $k$ )*

- (2) *the functor  $\Omega$  preserves the direct product:*

$$\Omega(X \times Y) = \Omega(X) \times \Omega(Y)$$

$$\Omega(f \times g) = \Omega(f) \times \Omega(g)$$

$$s_{X \times Y} = s_X \times s_Y, t_{X \times Y} = t_X \times t_Y$$

$$(\gamma_1 \times \gamma_2) \cdot_X (\gamma'_1 \times \gamma'_2) = (\gamma_1 \cdot_X \gamma'_1) \times (\gamma_2 \cdot_X \gamma'_2)$$

(the direct product is taken in the category of  $\mathfrak{S}ets$ .)

- (3) *unique path-lifting property*: if  $p \in \text{Hom}_{\mathfrak{E}}(X, Y)$  is an étale morphism of algebraic varieties, then for any points  $x, y \in \text{Pt}(X)$  the map

$$\Omega(p) : \bigcup_{y \in \text{Pt}(X)} \Omega_{x,y}(X) \rightarrow \bigcup_{z \in \text{Pt}(Y)} \Omega_{p(x),z}(Y)$$

is a bijection.

2.3.3. *Universality of the fundamental groupoid functor.* Let

$$\mathfrak{E} \subset \mathbb{V}, \text{Ob } \mathfrak{E} = \{E^n : n \geq 0\}, \text{Mor}_{\mathfrak{E}}(X, Y) = \text{Mor}_{\text{Var}/K}(X, Y)$$

be the full subcategory of the category of varieties whose objects are the Cartesian powers of  $E$  including  $E^0 = 0$  a variety consisting of the single point 0. By the definition of a full subcategory, the morphisms of  $\mathfrak{E}$  are morphisms of varieties between the objects of  $\mathfrak{E}$ .

The Galois group  $\text{Gal}(\overline{\mathbb{Q}}/k)$  acts on the category  $\mathfrak{E}$ ; the action leaves the objects invariant but permutes the morphisms. Recall we assume that all endomorphisms of  $E$  preserving  $0 \in E(k)$  are defined over its field  $k$  of definition.

Recall a groupoid  $\Omega(E)$  is *connected* iff for every  $x, y \in \text{Pt}(E)$  there exists  $\gamma \in \Omega(E)$  “going from point  $x$  to point  $y$ ”, i.e.  $x = s(\gamma), y = t(\gamma)$ . The set of such paths satisfying  $x = s(\gamma), y = t(\gamma)$  is denoted by  $\Omega_{x,y}(E)$ .

Using the notion of an abstract fundamental groupoid functor, we restate Proposition 2; then we show it is essentially equivalent to Proposition 1. The proof basically reconstructs the “universal covering space”  $V$  as the set of all paths  $\bigcup_{* \in E(\overline{\mathbb{Q}})} \Omega_{0,*}(E)$  leaving a particular point; functoriality of  $\underline{\Omega}$  allows us to define  $\text{End} E$ -module structure on  $V$ ; the unique path-lifting property of  $\Omega$  ensures unique divisibility.

**Proposition 2** (Universality of fundamental group functor). *Assume that  $E$  is an elliptic curve defined over a number field  $k$  and does not have complex multiplication. Let  $\mathfrak{E}$  be the full subcategory of Cartesian powers of  $E$  as above.*

*Let  $\underline{\Omega}$  be an abstract fundamental groupoid  $\overline{\mathbb{Q}}$ -valued functor on category  $\mathfrak{E}$ . Assume*

- (1)  $\underline{\Omega}(E)$  is a connected groupoid
- (2) there is an isomorphism

$$\Omega_{0,0}(E) \cong \mathbb{Z}^2 \text{ as } \text{End} E\text{-modules.}$$

Then there exists an automorphism  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that the 2-functors  $\underline{\Omega}, \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma : \mathfrak{E} \rightarrow \mathfrak{Sets}$  are naturally equivalent (as 2-functors),

$$\begin{array}{ccc} \underline{\Omega} & \cong & \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma \\ \mathfrak{E} & \xrightarrow{\quad \underline{\Omega} \quad} & \mathfrak{Sets} \\ \sigma \downarrow & & \downarrow \text{natural transformation} \\ \mathfrak{E} & \xrightarrow{\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}} & \mathfrak{Sets} \end{array}$$

If  $E$  has complex multiplication, then there exists finitely many abstract fundamental groupoid  $\overline{\mathbb{Q}}$ -valued functors  $\mathbb{F}_1, \dots, \mathbb{F}_n : \mathfrak{E} \rightarrow \mathfrak{Groupoids}$  such that for any other abstract fundamental groupoid  $\overline{\mathbb{Q}}$ -valued functor  $\underline{\Omega}$  there exists a Galois automorphism  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  and a number  $0 < i < n + 1$  such that  $\underline{\Omega}$  and  $\mathbb{F}_i \circ \sigma$  are naturally equivalent.

Recall  $\text{Pt}_{\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}}(V) = V(\overline{\mathbb{Q}})$  and  $\Omega_{\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}}(V) = \{\gamma \in \pi_1^{\text{top}}(V(\mathbb{C})) : s(\gamma), t(\gamma) \in V(\overline{\mathbb{Q}})\}$ .

The condition in the definition of an abstract fundamental groupoid functor are somewhat reminiscent of the conditions defining the scheme-theoretic algebraic fundamental group  $\pi_1^{\text{alg}}$  [SGA4 $\frac{1}{2}$ ]; however, there  $\pi_1^{\text{alg}}$  takes values in the category of *profinite* groups, in particular  $\pi_1^{\text{alg}}(\mathbb{C}^*, 1) = \hat{\mathbb{Z}}$ ,  $\pi_1^{\text{alg}}(E(\mathbb{C}), 0) = \hat{\mathbb{Z}}^2$ .

It is natural to consider whether a path lies in an algebraic subvariety, so if  $\underline{\Omega}$  is provide a useful notion of a path on  $E(\mathbb{C})$ , the 2-functor  $\underline{\Omega}$  restricted to  $\mathfrak{E}$  should be able to express when (a representative of the homotopy class of) a path lies in an *arbitrary* algebraic subvariety. This is indeed the case:

**Remark 3** (Recovering  $\Omega(Z)$  for arbitrary closed subvariety  $Z$  of  $E^n$ ). *The information contained in the functor  $\pi_1^{\text{top}}|_{\mathfrak{E}}$  restricted to the full subcategory  $\mathfrak{E}$  of Cartesian powers of an elliptic curve is enough to determine whether a path lies in a closed subvariety. The key fact here is that for a normal subgroup  $H \triangleleft \pi_1(E^n(\mathbb{C}))$ , there exists an H-Shafarevich morphism  $\text{Sh}_H : E^n \rightarrow E^m$  such that for an arbitrary irreducible  $Z \subset E^n(\mathbb{C})$ , it holds  $Z \subset \ker f$  iff the image  $\text{Im}[\pi_1(\hat{Z}, z) \rightarrow \pi_1(E^n(\mathbb{C}), z)]$  has a finite index subgroup contained in  $H$ .*

### 3. EXTENSIONS OF $E(\overline{\mathbb{Q}})$ BY $\Lambda$

In this section we state and prove Proposition 2; the proof is a model theoretic argument based on Kummer theory and the description of the image of Galois representation on Tate module  $T(E)$ . However, we



tried to be very explicit and have avoided any model-theoretic terminology in the exposition of the proof. The only model theory left in the proof is in the level of motivations; as was noted earlier, those model-theoretic motivations are useful to try and generalise the formulation to other varieties and contexts in general.

**3.1. Transitivity of  $\text{Gal}(K/\mathbb{Q})$  action on the uniquely divisible  $\text{End } E$ -module extensions of  $E(\overline{\mathbb{Q}})$  by  $\Lambda$ .** To fix notations for the proof, we restate Proposition 1 in an expanded form.

**Proposition 4.** *Let  $E(\mathbb{C}), \Lambda$  be as above.*

- (1) *There exists a uniquely divisible  $\text{End } E$ -module  $V$  and a short exact sequence of  $\text{End } E$ -modules*

$$0 \longrightarrow \Lambda \longrightarrow V \longrightarrow E(\overline{\mathbb{Q}}) \longrightarrow 0.$$

- (2) (a) *If  $E$  has no complex multiplication and  $\text{End } E = \mathbb{Z}$ , then for any uniquely divisible  $\mathbb{Z}$ -module extensions  $W, V$  of  $E(\overline{\mathbb{Q}})$  by  $\Lambda$ , fitting into the short exact sequences as above, there exist a commutative diagram:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda & \longrightarrow & V & \xrightarrow{\varphi} & E(\overline{\mathbb{Q}}) \longrightarrow 0 \\ & & \exists h|_{\Lambda} \downarrow & & \exists h \in \text{Hom}(V, W) \downarrow & & \exists \sigma \in \text{Gal}(\overline{\mathbb{Q}}/k) \downarrow \\ 0 & \longrightarrow & \Lambda & \longrightarrow & W & \xrightarrow{\psi} & E(\overline{\mathbb{Q}}) \longrightarrow 0 \end{array}$$

- (b) *If  $\text{End } E \neq \mathbb{Z}$ , then there exist finitely many uniquely divisible  $\text{End } E$ -module extensions  $W_1, \dots, W_n$  of  $E(\overline{\mathbb{Q}})$  by  $\Lambda$ , fitting into the short exact sequences as above, such that for any uniquely divisible  $\text{End } E$ -module extension  $V$  there exist a commutative diagram:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda & \longrightarrow & V & \xrightarrow{\varphi} & E(\overline{\mathbb{Q}}) \longrightarrow 0 \\ & & \exists h|_{\Lambda} \downarrow & & \exists h \in \text{Hom}(V, W) \downarrow & & \exists \sigma \in \text{Gal}(\overline{\mathbb{Q}}/k) \downarrow \\ 0 & \longrightarrow & \Lambda & \longrightarrow & W_i & \xrightarrow{\psi_i} & E(\overline{\mathbb{Q}}) \longrightarrow 0 \end{array}$$

*Proof.* By assumption there is a covering

$$0 \longrightarrow \Lambda \longrightarrow \mathbb{C} \xrightarrow{p} E(\mathbb{C}) \longrightarrow 0$$

The endomorphisms act on the complex plane  $\mathbb{C}$  by multiplication by complex numbers, and so in particular  $\mathbb{C}$  is a uniquely divisible  $\text{End } E$ -module. The set  $E(\overline{\mathbb{Q}})$  of points of  $E$  over an algebraically closed subfield is closed under addition and  $\text{End } E$ -multiplication i.e. is an  $\text{End } E$ -submodule, necessarily uniquely divisible; so is then  $V = p^{-1}(E(\overline{\mathbb{Q}}))$ ; take that to obtain a short exact sequence as above. This proves (1).

To prove (2), we use the Kummer theory of elliptic curves and the Serre's results about the image of Galois representation on the torsion points. Essentially, what we need is that all the restrictions on the image of the Galois action on the Tate module  $\prod_l T(E)$  come from the geometry of  $E$  and in this case, are linear. That is exactly what is provided by the Kummer theory and other results we use.

Pick a maximal linearly independent set  $v_0, v_1, v_2, \dots \in V$ ; let  $V_n = \text{End}E v_0 + \dots + \text{End}E v_n$  be the submodule generated by  $v_0, \dots, v_n$ , and let  $\mathbb{Q}V_n = (\text{End}E)^{-1}V_n = \{v : \exists N \in \mathbb{N}(Nv \in V_n)\}$  be its divisible closure. We construct by induction a partial  $\text{End}E$ -module linear map  $h_n : \mathbb{Q}V_n \rightarrow W_i$  inducing a partial Galois map  $\sigma_n : \varphi(\mathbb{Q}V) \rightarrow E(\overline{\mathbb{Q}})$  so that  $h = \cup h_n$  is an isomorphism of  $V$  and  $W_i$ ; then the construction implies  $\sigma = \cup \sigma_n$  is a total Galois map on  $E(\overline{\mathbb{Q}})$  to  $E(\overline{\mathbb{Q}})$ , and thus there is a commutative diagram as above.

For a point  $a \in E(\mathbb{C})$ , let us call a *compatible sequence of division points starting at  $a$*  a sequence  $(a^{(i)})_{i \in \mathbb{N}}$  satisfying  $a^{(i)} = ja^{(ij)}$ ,  $i, j \in \mathbb{N}$ . Compatible sequences of division points starting at  $0 \in E(\overline{\mathbb{Q}})$  form a *Tate module*  $T(E)$ . A compatible sequence of  *$l$ -primary* division points is a subsequence  $(a^{(l^j)})_{j \in \mathbb{N}}$  of a compatible sequence of division points; such sequences of  $l$ -primary points starting at 0 form  *$l$ -adic Tate module*  $T_l(E)$ .

### 3.2. The image of Galois representations on Tate module $T_l(E)$ .

**3.2.1. Base of induction.** Choose  $v_0 \in \Lambda = \ker \varphi$ ; then  $\mathbb{Q}V_0 = \varphi^{-1}(E_{\text{tors}})$ , where  $E_{\text{tors}} = \{x \in E(\overline{\mathbb{Q}}) : nx = 0 \text{ for some } n \in \mathbb{N}\}$  is the set of *torsion* points of  $E(\overline{\mathbb{Q}})$ .

As the base step of the induction, in this subsection we construct a commutative diagram:

$$(1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \Lambda & \longrightarrow & V_0 & \xrightarrow{\varphi} & E_{\text{tors}} \longrightarrow 0 \\ & & \downarrow h_0 & & \downarrow h_0 \in \text{Hom}(V_0, W) & & \downarrow \sigma \in \text{Gal}(\overline{\mathbb{Q}}/k) \\ 0 & \longrightarrow & \Lambda & \longrightarrow & W & \xrightarrow{\psi \circ \tau_i} & E_{\text{tors}} \longrightarrow 0 \end{array}$$

where  $\tau_1, \dots, \tau_n$  is some fixed finite collection of  $\text{End}E$ -endomorphisms of  $E_{\text{tors}}$  independent of  $V, W$ . For curves without complex multiplication, there is no need to consider  $\tau_i$ 's, i.e.  $n = 1$ ,  $\tau_1 = \text{id}$ .

Note that when  $E$  has complex multiplication, the diagram above is somewhat reminiscent of a diagram appearing in the main theorem of complex multiplication [Shi71, Chapter 5, Theorem 5.4], also Lang [Lan83]. Global class field theory provides an explicit description on

the image of Galois action in  $\text{Aut}(E_{\text{tors}})$  in terms of the action of the ring of idèles on some lattices in the universal covering space.

**3.2.2.  $E$  has complex multiplication.** Pick an arbitrary isomorphism  $h_0 : \Lambda \rightarrow \Lambda$  and extend it uniquely to  $h_0 : V_0 \rightarrow W$ ; we may do so by unique divisibility of  $V$  and  $W$ . Define an  $\text{End}E$ -morphism  $\tau : E_{\text{tors}} \rightarrow E_{\text{tors}}$  by  $\tau(x) = \psi \circ h_0 \circ \varphi^{-1}(x)$ . The calculation  $\psi \circ h_0(y + \Lambda) = \psi \circ h_0(y)$  shows it is well-defined; linearity of  $\tau$  follows from that of  $\varphi, h_0$  and  $\psi$ .

Ideally we would like to be able to choose  $h_0 : \Lambda \rightarrow \Lambda$  so that  $\tau = \tau_h : E_{\text{tors}} \rightarrow E_{\text{tors}}$  is induced by a Galois automorphism. Here we prove a weaker statement below.

Denote  $\mathcal{O} = \text{End}E$  and  $E[n] = \{x \in E(\overline{\mathbb{Q}}) : nx = 0\}$  the  $n$ -torsion of  $E$ . The set  $E[n]$  is a free 1-dimensional  $\mathcal{O}/n\mathcal{O}$ -module ([Lan78, Ch.8§15, Fact 1]), and  $\text{Aut}_{\mathcal{O}}(E[n]) = \text{Aut}_{\mathcal{O}/n\mathcal{O}\text{-mod}}(E[n]) \cong (\mathcal{O}/n\mathcal{O})^*$ , where  $(\mathcal{O}/n\mathcal{O})^*$  denotes the group of invertible elements in  $(\mathcal{O}/n\mathcal{O})^*$ .

An  $\mathcal{O}$ -automorphism of  $E_{\text{tors}}$  is given by a compatible system of  $\mathcal{O}$ -automorphisms of  $E[n]$ ,  $n > 0$ ; thus we see that there is an action of  $\hat{\mathcal{O}} = \varprojlim_n \mathcal{O}/n\mathcal{O}$  on  $E_{\text{tors}}$  as an  $\text{End}E$ -module; the fact that  $\text{Aut}_{\mathcal{O}}(E[n]) = \text{Aut}_{\mathcal{O}/n\mathcal{O}\text{-mod}}(E[n]) \cong (\mathcal{O}/n\mathcal{O})^*$  implies that  $\text{Aut}_{\mathcal{O}}(E_{\text{tors}}) \cong \hat{\mathcal{O}}$ .

Now we refer to a consequence of the main theorem of complex multiplication, namely that, in notation of [Lan78, Ch.8§15, Fact 2], the image of Galois group

$$G_K = \text{Gal}(K(E_{\text{tors}}) : K) \rightarrow \prod ((\text{End}E)_l)^*$$

is open of finite index, i.e.  $\text{Im} G_K$  is a finite index subgroup of  $\hat{\mathcal{O}}^* = \prod_l \mathcal{O}_l^*$ . Choose  $\tau_1, \dots, \tau_n$  to be representatives of conjugacy classes  $\mathcal{O}^*/\text{Im} G_K$ ; we then have that for some  $i$   $\tau_i \tau = \sigma \in \text{Im} G_K$ ; this choice of  $h_0, \sigma = \tau_i \tau$  makes the diagram (1) commutative, as required.

**3.2.3.  $E$  does not have complex multiplication.** Assume that  $E$  does not have complex multiplication, i.e.  $\text{End} E \cong \mathbb{Z}$ ; identify  $\text{End}E = \mathbb{Z}$ ,  $T(E) = \hat{\mathbb{Z}}^2$ , and  $\text{Aut}(T(E)) = \text{SL}_2(\hat{\mathbb{Z}})$ . The maps  $\varphi : V \rightarrow E(\overline{\mathbb{Q}})$ ,  $\psi : W \rightarrow E(\overline{\mathbb{Q}})$  define embeddings  $\iota_\varphi : \Lambda \rightarrow T(E)$ ,  $\iota_\psi : \Lambda \rightarrow T(E)$  by

$$\iota_\varphi : \lambda \mapsto (\varphi(\lambda/j))_{j \in \mathbb{N}},$$

$$\iota_\psi : \lambda \mapsto (\psi(\lambda/j))_{j \in \mathbb{N}}.$$

The images  $\iota_\varphi(\Lambda), \iota_\psi(\Lambda)$  of the both maps are dense in  $T(E)$  due to the surjectivity of  $\varphi, \psi : \mathbb{Q}\Lambda \rightarrow E_{\text{tors}}$ .

Take a pair of elements  $\lambda_0, \lambda_1 \in \ker \varphi \cong \Lambda$  generating  $\Lambda$  as an Abelian group; we want to find  $\lambda'_0, \lambda'_1 \in \ker \psi \cong \Lambda$  and  $\sigma = \sigma_0 \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that

$$\sigma \iota_\varphi(\lambda_0) = \iota_\psi(\lambda'_0),$$

$$\sigma \iota_\varphi(\lambda_1) = \iota_\psi(\lambda'_1).$$

Under identification  $\ker \varphi = \mathbb{Z}^2$ , since vectors  $\lambda_0, \lambda_1 \in \mathbb{Z}^2$  generate lattice  $\mathbb{Z}^2$ , it holds that  $\det(\lambda_0, \lambda_1) = 1$ . That implies that  $\det(\iota_\varphi(\lambda_0), \iota_\varphi(\lambda_1))$  is a unit in  $\hat{\mathbb{Z}} = \prod_l \mathbb{Z}_l$ . Similarly  $\det(\iota_\psi(\lambda'_0), \iota_\psi(\lambda'_1))$  has to be a unit in  $\hat{\mathbb{Z}}$ .

By [Ber88, Theorem 3], the image of the Galois group  $\text{Gal}(\overline{\mathbb{Q}}/k)$  in the automorphism group  $\text{Aut}(T(E)) = \text{SL}_2(\hat{\mathbb{Z}})$  contains an open subgroup  $\text{SL}_2(N\hat{\mathbb{Z}}) = \ker \left( \text{SL}_2(\hat{\mathbb{Z}}) \rightarrow \text{SL}_2(\mathbb{Z}/N\mathbb{Z}) \right)$ , for some  $N \in \mathbb{N}$  large enough. Now, there is  $L \in \text{SL}_2(N\hat{\mathbb{Z}})$  such that  $L(u_0) = u'_0$  and  $L(u_1) = u'_1$  iff  $\det(u_0, u_1) = \det(u'_0, u'_1)$  and  $u_0 \in u'_0 + N\hat{\mathbb{Z}}$ ,  $u_1 = u'_1 + N\hat{\mathbb{Z}}$ . Thus it is enough to take  $\mathbb{Z}$ -linearly independent  $\lambda'_0, \lambda'_1$  such that  $\psi(\lambda'_0/N) = \varphi(\lambda_0/N)$  and  $\psi(\lambda'_1/N) = \varphi(\lambda_1/N)$ . Necessarily then  $\det(\iota_\psi(\lambda'_0), \iota_\psi(\lambda'_1))$  is a unit. This implies that  $\det(\lambda'_0, \lambda'_1) = 1$  and thus,  $\lambda'_0, \lambda'_1$  generate  $\mathbb{Z}^2$ . The latter statement is independent of the identification  $\ker \psi = \mathbb{Z}^2$ , and this concludes the proof in the case of no complex multiplication.

### 3.3. Kummer theory.

**3.3.1. The main statement of the Kummer theory for an elliptic curve.** Let us state the main lemma in a form convenient to us to make an inductive process; that is a form natural from the model-theoretic point of view and corresponds to the property of *atomicity of certain formulae over the kernel*.

**Lemma 5** (Kummer theory for an elliptic curve). *Let  $E$  be an elliptic curve defined over a number field  $k$ . Let  $a_1, \dots, a_n \in E(\overline{\mathbb{Q}})$  be a sequence of points linearly independent over  $\text{End} E$ . Then there exists  $N \in \mathbb{Z}$  such that any two compatible sequences  $(a_1^{(i)}, \dots, a_n^{(i)})_{i \in \mathbb{N}}$ ,  $(b_1^{(i)}, \dots, b_n^{(i)})_{i \in \mathbb{N}}$  of division points in  $E(\overline{\mathbb{Q}})$  starting at  $a_1, \dots, a_n$  and such that  $a_1^{(N)} = b_1^{(N)}, \dots, a_n^{(N)} = b_n^{(N)}$ , are  $\text{Gal}(\overline{\mathbb{Q}}/k)$ -conjugated by  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$ ,*

$$\sigma(a_i^{(j)}) = b_i^{(j)}, \text{ for all } 0 \leq i \leq n, j \in \mathbb{N}.$$

*Proof of Lemma.* See Bashmakov [Bas72] for original results for elliptic curves; see [Rib79, JR87, Rib87] for Kummer theory of Abelian varieties, and see [Ber88] for a summary of results of Kummer theory of Abelian varieties; we quote [Ber88, Theorem 2] from that paper.

We now introduce the notations of Bertrand [Ber88, Theorem 2]. Let  $G = A \times L$  be a product of an Abelian variety by a torus  $L$  so that after a finite extension of  $k$  it satisfies Poincaré's complete

reducibility theorem (as a variety *over*  $k$ ). Let  $l$  denote a prime. Let  $G_{l^\infty} = \{x \in G(\bar{k}) : \exists n \, l^n x = 0\}$  be the  $l^\infty$ -torsion of  $G$ . For a point  $P \in G(k)$ , let  $G_P$  be the smallest algebraic subgroup of  $G$  containing  $P$ , i.e. the Zariski closure of subgroup  $\mathbb{Z}P$  of  $G$ , and let  $G_P^\circ$  be its connected component through the origin, and finally let

$$\xi_{l^\infty}(P) : \text{Gal}(\bar{k}/k(G_{l^\infty}, P)) \longrightarrow T_{l^\infty}(A \times L)$$

$$\sigma \longmapsto \sigma(P_{l^\infty}) - P_{l^\infty}, \text{ for some } P_{l^\infty} \in T_{l^\infty}(A \times L).$$

It can be checked by direct computation the map  $\xi_{l^\infty}(P) : \text{Gal}(\bar{k}/k(G_{l^\infty}, P)) \rightarrow T_{l^\infty}(A \times L)$  does not depend on the choice of  $P_{l^\infty} \in T_{l^\infty}(A \times L)$ .

Let  $T(G_P^\circ)$  be the sequences of  $T(E)$  consisting of elements of  $G_P^\circ$ ; then, according to [Ber88, Theorem 2], the image of  $\xi_\infty(P) = \prod_l \xi_{l^\infty}$  has finite index in  $T(G_P^\circ) = \prod_l T_{l^\infty}(A \times L)$ , i.e. the image contains  $NT(G_P^\circ)$ , for some natural number  $N \in \mathbb{N}$  large enough.

We claim that if we take  $G = E^n$ ,  $P = (a_1, \dots, a_n) \in E^n(\overline{\mathbb{Q}})$ , then  $N$  above is  $N$  required in Lemma. By the result cited above, it is enough to prove that if  $a_1, \dots, a_n \in E(\overline{\mathbb{Q}})$  are  $\text{End}E$ -linearly independent, then  $G_P^\circ = E^n$ .

An algebraic subgroup of  $E^n$  is necessarily an Abelian subvariety; by the Corollary of Poincaré's complete reducibility theorem (Lemma 6 below) if  $\dim G_P^\circ \neq E^n$ , then for some  $m \in \mathbb{N}$ , the connected component of  $G_P^\circ/m$  lies in the kernel of some non-trivial morphism  $f : E^n \rightarrow E^m$ . The kernel of  $f$  satisfies the  $\text{End}E$ -linear relation corresponding to  $f : E^n(\bar{k}) \rightarrow E^m(\bar{k})$  as a morphism of  $\text{End}E$ -modules. Since by assumption  $P$  satisfies no  $\text{End}E$ -linear relation, this is a contradiction, and  $G_P^\circ = E^n$ .

**3.3.2. Any Abelian subvariety is a connected component of the kernel of a morphism.** The following Lemma has been just used to relate the more geometric formulation of Bertrand [Ber88] and the more explicit statement of Lemma 5 following Bashmakov [Bas72] and Ribet [Rib79] specific to Abelian varieties.

**Lemma 6** (Corollary of Poincaré's complete reducibility theorem). *Let  $B \subset E^n$  be an irreducible Abelian subvariety. Then there is a morphism  $f : E^n \rightarrow E^m$  such that  $B$  is a connected component of  $\ker f$ .*

*In particular, this implies that an Abelian subvariety  $B \neq E^n$  satisfies a non-trivial  $\text{End}E$ -linear relation.*

*Proof.* By Poincaré's reducibility theorem, there exist an Abelian subvariety  $B' \subset E^n$  such that  $B(\bar{k}) + B'(\bar{k}) = E^n(\bar{k})$  and the intersection  $B(\bar{k}) \cap B'(\bar{k})$  is finite. This implies there is an isogeny  $f : B \times B' \rightarrow E^n$ ,  $(x, y) \mapsto (x + y)$ , and  $f(B) = B$ . There exists another isogeny

$f' : E^n \rightarrow B \times B'$  such that  $f' \circ f = [m]$  is a multiplication by a natural number  $m \in \mathbb{N}$ ; let  $B_m$  and  $B'_m$  be the connected components of  $B/m \subset E^n$  and  $B'/m \subset E^n$ , respectively, passing through the origin. Then we know that  $mB_m = ff'(B_m) = B$ ,  $mB'_m = ff'(B'_m) = B'$  but also  $f'(B_m)$  and  $f'(B'_m)$  do not intersect within  $B \times B'$ ; to conclude,  $B_m, B'_m \subset E^n$  and  $B_m \cap B'_m = \emptyset$  and  $B_m + B'_m = E^n$ . Let  $p_2 : B \times B' \rightarrow B'$  be the projection on the second coordinate; since  $0$  is a connected component of  $\ker f$ , it implies that  $B_m$  is a connected component of kernel of  $f \circ p_2 \circ f' : E^n \rightarrow E^n$  and  $B$  is a connected component of  $f \circ p_2 \circ f' \circ [m] : E^n \rightarrow E^n$ .  $\square$

This concludes the proof of the lemma 5 stating the Kummer theory for an elliptic curve.  $\square$

**3.3.3. An inductive argument based on Kummer theory.** Assume now that we are on inductive step  $n-1$ , i.e. we have defined an  $\text{End}E$ -linear map  $h_{n-1} : \mathbb{Q}V_{n-1} \rightarrow W$  and  $\sigma_{n-1} \in \text{Gal}(\overline{\mathbb{Q}}/k)$ ,  $h_{n-1}(v_i) = w_i, 0 < i < n$  such that  $\psi(h_{n-1}(v)) = \sigma_{n-1}\varphi(v)$  for every  $v \in \mathbb{Q}V_{n-1}$ . Consider a compatible system  $(\varphi(v_1/j))_j, \dots, (\varphi(v_n/j))_j, j \in \mathbb{N}$  of division points in  $E(\overline{\mathbb{Q}})$ , and take  $N$  as in Kummer theory Lemma 5. By the induction hypothesis we have  $\sigma_{n-1}\varphi(v_1/j) = \psi(v_1/j), \dots, \sigma_{n-1}\varphi(v_{n-1}/j) = \psi(v_{n-1}/j)$  for any  $j$ . Choose  $w_n \in W$  such that  $\sigma_{n-1}\varphi(v_n/N) = \psi(w_n/N)$ ; that is possible by surjectivity of  $\psi : W \rightarrow E(\overline{\mathbb{Q}})$ . By Kummer theory lemma, for  $N$  large enough, there exists  $\sigma' \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that  $\sigma'\sigma_{n-1}\varphi(v_1/j) = \psi(v_1/j), \dots, \sigma'\sigma_{n-1}\varphi(v_{n-1}/j) = \psi(v_{n-1}/j)$ , and  $\sigma'\sigma_{n-1}\varphi(v_n/j) = \psi(w_n/j)$ ; let  $\sigma_n = \sigma'\sigma_{n-1}$ . By construction we have that  $\sigma_n|_{\varphi(\mathbb{Q}V_{n-1})} = \sigma_{n-1}|_{\varphi(\mathbb{Q}V_{n-1})}$  and  $\sigma_n\varphi(v_i/j) = \psi(v_i/j), 0 < i < n+1$ . This implies  $\sigma_n\varphi(v) = \psi(v)$  for arbitrary  $v \in \mathbb{Q}V_n$ , thereby completing the induction step.

After countably many steps we construct a total  $\text{End}E$ -linear map  $h = \cup h_n : V \rightarrow W$  and  $\sigma : \varphi(V) \rightarrow E(\overline{\mathbb{Q}})$ . Since  $\varphi(V) = E(\overline{\mathbb{Q}})$ , the Galois map  $\sigma$  is defined on the whole of  $E(\overline{\mathbb{Q}})$ . Since Galois map  $\sigma$  is surjective, this implies  $h : V \rightarrow W$  is surjective, too. This completes the proof of Proposition 1.

The last argument could in fact have been avoided by a little more careful inductive construction of  $h$ : instead of always choosing  $w_n$  to match  $\varphi(v_n)$  we could have on odd steps pick an arbitrary  $w_n$  and then chosen  $v_n$  so that  $\sigma_n(\varphi(v_n)) = \psi(w_n)$  while on even steps preserving the old behaviour. It is very easy to force surjectivity of the constructed map  $h : V \rightarrow W$  this way; it is a very common argument in model theory called “a back-and-forth argument”.  $\square$

### 3.4. Concluding Remarks.

### 3.4.1. Image of Galois representation.

**Remark 7.** *Note that for the arguments in §3.2 it is essential that the image of the Galois action is as large as possible subject to linear dependencies. However, this is something specific to elliptic curves and false for higher dimensional Abelian varieties: one needs to take care of a symplectic form. This observation shows that the straightforward generalisation to higher dimensional Abelian varieties is false. See [Gav05, IV§7] for a discussion of a way to do that.*

3.4.2. *Kummer theory.* Remark 7 says that in higher dimensions, the base of the induction breaks down due to additional restrictions on the image of the Galois action. This does not happen in the later steps of the induction process based on Kummer theory.

**Remark 8** (Generalisations of Kummer theory argument). *Since Kummer theory is known in much larger generality, say for a product of arbitrary Abelian varieties, complex tori  $\mathbb{C}^*$  and complex lines  $\mathbb{C}$  ([Ber88]), it seems straightforward to generalise the Kummer theory argument above to such a product  $A$ . Thus, one would prove that if there exists  $h_0 : \ker \varphi \rightarrow \ker \psi$  and a Galois map  $\sigma_0 : E(k(T(E))) \rightarrow E(k(T(E)))$  in  $\text{Gal}(k(T(E))/k)$  such that  $\varphi \circ \sigma_0 = h_0 \circ \psi$ , then there exists  $h : V \rightarrow W$  and  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  making the diagram (2a) commute. That is, a morphism between kernel from  $\Lambda \subset V$  to  $\Lambda \subset W$  extends to a morphism on the whole of  $V$ . Model-theoretically, this means that the types of the points of universal covering space lying over algebraic points  $A(\overline{\mathbb{Q}})$  is atomic over the kernel in the linear language.*

**Remark 9** (Failure of Kummer theory for extensions of Abelian varieties by tori). *According to Ribet [Rib87, JR87], Kummer theory may fail for non-trivial extensions of Abelian varieties by  $(\mathbb{C}^*)^n$  due to “existence of an additional morphism”; he gives a motivic interpretation in [Rib87]. It is natural to ask if an analogous argument could still be carried despite the failure of Kummer theory. To state a correct conjecture, we may need to use more general considerations of [Gav05].*

## 4. FUNDAMENTAL GROUPOID FUNCTOR

In this section we derive Proposition 2 from Proposition 1 by carrying out a formal counterpart of a natural topological construction of a universal covering space.

**4.1. Uniqueness of extensions implies the universality of the fundamental groupoid functor.** In this § we show that Proposition 1 implies Proposition 2. We do so by explicitly constructing an

extension as in Proposition 1 from a 2-functor as in Proposition 2, and then showing that the equivalence of constructed extensions implies the equivalence of 2-functors.

The construction is a formalisation of the geometric observation that the universal covering space with a basepoint can be canonically identified with the set of homotopy classes of paths leaving the basepoint of the base. The identification is via the unique path-lifting property of the covering map, and depends only on the choice of a basepoint in the universal covering space. In case of the universal covering space  $\mathbb{C}$  of  $E(\mathbb{C})$ , the correspondence is given by the *period map*

$$\gamma \longmapsto \int_{\gamma} dz.$$

4.1.1. *Recovering  $V$  as the universal covering space from the 2-functor  $\underline{\Omega}$ .* Let  $\underline{\Omega} = (\text{Pt}, \Omega, s, t, \cdot)$  be a 2-functor from  $\mathfrak{E}$  to  $\mathfrak{Sets}$  satisfying the conditions of Definition 1. We want to construct an extension

$$0 \longrightarrow \mathbb{Z}^2 \longrightarrow V_{\underline{\Omega}} \xrightarrow{\varphi} E(\overline{\mathbb{Q}}) \longrightarrow 0$$

of  $\text{End}E$ -modules. The topological intuition above suggests that we set

$$V = V_{\underline{\Omega}} = \bigcup_{y \in \text{Pt}(E)} \Omega_{0,y}(E) = \{\gamma \in \Omega(E) : s(\gamma) = 0\} \text{ (disjoint union),}$$

$$\varphi(\gamma) = t(\gamma)$$

where  $0 \in E(k)$  is the zero point of the elliptic curve  $E$ .

4.1.2. *Abelian group structure on  $\Omega(E)$ .* The functoriality of  $\underline{\Omega}$  transfers  $\text{End}E$ -module structure on  $E(\overline{\mathbb{Q}})$  to that on  $V$ ; namely, let us check that the maps  $\Omega(f)$ ,  $f \in \text{End}E$ , and  $\Omega(m)$ , where  $m : E \times E \rightarrow E$  is the morphism of addition on  $E$ , define  $\text{End}E$ -module structure on  $V$ , or rather that their restriction to  $V$  does.

By assumption the functor  $\underline{\Omega}$  preserves direct products so  $\Omega(E \times E) = \Omega(E) \times \Omega(E)$ , and thus there is a map

$$\Omega(m) : \Omega(E) \times \Omega(E) \rightarrow \Omega(E).$$

Maps  $s, t$  are natural transformations of  $\Omega$  to  $\text{Pt}$  (as functors to  $\mathfrak{Sets}$ ) and so  $s \circ \Omega(m) = \text{Pt}(m) \circ s$ ,  $t \circ \Omega(m) = \text{Pt}(m) \circ t$  is the map of addition on end-points. Therefore,

$$\Omega(m)(\Omega_{x,y}(E) \times \Omega_{v,w}(E)) \subset \Omega_{x+v, y+w}(E),$$

and in particular

$$\Omega(m)(\Omega_{0,y}(E) \times \Omega_{0,z}(E)) \subset \Omega_{0, y+z}(E).$$



Thus  $\Omega(m) : V \times V \rightarrow V$  gives us a binary operation. It is straightforward to show that the preservation of direct product and functoriality implies that  $\Omega(m)$  makes  $\Omega(E)$  into an Abelian group. Let us check this.

By definition, associativity of  $m : E \times E \rightarrow E$  means that  $m \circ (m \times \text{id}_E) = m \circ (\text{id}_E \times m) : E \times E \times E \rightarrow E$ ; by preservation of direct product this implies  $\Omega(m) \circ (\Omega(m) \times \text{id}_E) = \Omega(m) \circ (\text{id}_E \times \Omega(m))$  and so  $\Omega(m)$  is associative. Similarly, commutativity of  $m$  means  $m \circ (\text{id}_1 \times \text{id}_2) = m \circ (\text{id}_2 \times \text{id}_1) : E \times E \rightarrow E$ ; that similarly implies the commutativity of  $\Omega(m)$ . In the language of morphisms, the existence of a zero for the additive law translates to the existence of a morphism  $0 : \{0\} \rightarrow E$  subject to the identifies:  $m \circ (\text{id} \times 0) = p_2 : \{0\} \times E \rightarrow E$  and  $m \circ (0 \times \text{id}) = p_1 : E \times \{0\} \rightarrow E$  corresponding to a commutative diagram:

$$\begin{array}{ccc} \{0\} \times E & \longrightarrow & E \times E \\ & \searrow & \swarrow \\ & E & \end{array}$$

Apply  $\Omega$  to get  $\Omega(m) \circ (\Omega(\text{id}) \times \Omega(0)) = \Omega(p_2) : \Omega(\{0\}) \times \Omega(E) \rightarrow \Omega(E)$  and  $\Omega(m) \circ (\Omega(0) \times \Omega(\text{id})) = \Omega(p_1) : \Omega(E) \times \Omega(\{0\}) \rightarrow \Omega(E)$ . Preservation of direct product implies  $\Omega(m) \circ (\text{id} \times \Omega(0)) = \Omega(p_2) : \Omega(\{0\}) \times \Omega(E) \rightarrow \Omega(E)$  and  $\Omega(m) \circ (\Omega(0) \times \text{id}) = \Omega(p_1) : \Omega(E) \times \Omega(\{0\}) \rightarrow \Omega(E)$ . This implies that  $\Omega(0)(\Omega(\{0\}))$  is a zero point in  $V$ .

Existence of (right) inverse corresponds to the existence of a morphism  $i : E \rightarrow E$  subject to the following commutative diagram:

$$\begin{array}{ccc} E & \longrightarrow & \{0\} \\ (\text{id}_E, i) \downarrow & & \downarrow 0 \\ E \times E & \xrightarrow{m} & E \end{array}$$

Again functoriality ensures that  $\Omega(i)$  satisfies a similar diagram, thus proving the existence of inverses.

The above checks that  $\Omega(m)$  is an associative commutative partial operation on  $\Omega(E)$  possessing a zero element and inverses; it is immediate to check  $V$  is closed under  $\Omega(m)$  and inverse  $\Omega(i)$ , and so is a group.

**4.1.3. Action of “fundamental group”  $\Omega_{0,0}(E)$  on  $V$  via concatenation and by  $\Omega(m)$ -multiplication.** Take a loop  $\lambda \in \Omega_{0,0}(E)$  and  $\gamma \in \Omega_{0,y}$ ; then both concatenation and  $\Omega(m)$ -product of  $\lambda$  and  $\gamma$  are well-defined; let us show that  $\lambda \cdot \gamma = \Omega(m)(\lambda \times \gamma)$ :

$$\Omega(m)(\lambda \times \gamma) = \Omega(m)(\lambda \cdot 0 \times 0 \cdot \gamma) = \Omega(m)(\lambda \times 0) \cdot \Omega(m)(0 \times \gamma) = \lambda \cdot \gamma.$$

The latter equality follows from the inverse element equality of morphisms  $m(\text{id} \times 0) = m(0 \times \text{id}) = \text{id}$ . In the classical example, this observation corresponds to the following calculation:

$$\int_{\gamma_1 \cdot \gamma_2} dz = \int_{\gamma_1} dz + \int_{\gamma_2} dz = \int_{\gamma_1 \gamma_2} dz.$$

Here  $\gamma_1 \cdot \gamma_2$  denotes the concatenation of the paths and  $\gamma_1 \gamma_2$  denotes the pointwise product of the paths.

4.1.4. *Divisibility of  $\text{End}E$ -module structure and path-lifting property.* Analogously, a morphism  $f \in \text{End}E$ ,  $\Omega(f) : \Omega(E) \rightarrow \Omega(E)$  defines a map  $\Omega(f) : \Omega(E) \rightarrow \Omega(E)$ . Arguments similar to the ones above allow us to prove that  $V$  is an  $\text{End}E$ -module with the operations defined above. Let us now prove that the path-lifting property implies that  $V$  is uniquely divisible. Indeed, we know that any isogeny  $f \in \text{End}E$  is étale ([Mum70]) and we may apply the path-lifting property to get a bijection

$$\Omega(f) : \bigcup_{0 \in \text{Pt}(E)} \Omega_{0,y}(E) \longrightarrow \bigcup_{z \in \text{Pt}(E)} \Omega_{0,z}(E).$$

That is, by definition of  $V$ , the map  $\Omega(f)$  is a bijection on  $V$ , and  $V$  is uniquely divisible as required.

Finally, to get a short exact sequence as in Proposition 1, set  $\varphi(w) = t(w)$ . Then naturality of target map  $t$  implies  $\varphi(w)$  is homomorphism of  $\text{End}E$ -modules; the connectivity of  $\Omega(E)$  implies that  $\varphi$  is surjective. The kernel  $\ker \varphi$  of  $\varphi : V \rightarrow E(\overline{\mathbb{Q}})$  is  $\Omega_{0,0}(E)$  and is isomorphic to  $\mathbb{Z}^2$  by assumption.

4.1.5. *Construction of a natural transformation  $h' : \underline{\Omega} \rightarrow \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma$ .* For notational convenience, let  $\underline{\Omega}' = (\text{Pt}, \Omega', s', t', \cdot')$  denote the functor  $\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}$ .

Let  $W \xrightarrow{\psi} E(\overline{\mathbb{Q}})$  be the  $\text{End}E$ -module extension constructed from  $\underline{\Omega}' = \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}}$  in the same way. Since  $V, W$  are both uniquely divisible extensions of  $E(\overline{\mathbb{Q}})$  by  $\mathbb{Z}^2$ , we may apply Proposition 1 to get an  $\text{End}E$ -linear map  $h : V \rightarrow W$  and  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that  $\sigma \circ \varphi = \psi \circ h$ .

We want to get a natural transformation  $h' : \underline{\Omega} \rightarrow \underline{\Omega}' \circ \sigma$ . Recall a natural transformation  $h' : \underline{\Omega} \rightarrow \underline{\Omega}' \circ \sigma$  is a family of maps (of sets with no further structure)  $h_A^\Omega : \Omega(A) \rightarrow \Omega' \circ \sigma(A)$  and  $h_A^{\text{Pt}} : \text{Pt}(A) \rightarrow \text{Pt}' \circ \sigma(A)$  satisfying certain compatibility conditions expressed by commutative diagrams (2),(3),(4).

Define  $h_A^{\text{Pt}} : A(\overline{\mathbb{Q}}) \rightarrow A(\overline{\mathbb{Q}})$  by  $h_A^{\text{Pt}}(x) = \sigma(x) \in \text{Pt}'(A) = (\sigma A)(\overline{\mathbb{Q}}) = A(\overline{\mathbb{Q}})$  to be the map induced by Galois automorphism  $\sigma : \overline{\mathbb{Q}} \rightarrow \overline{\mathbb{Q}}$ .

By assumption of connectivity of  $\Omega(E)$ , any  $\gamma \in \Omega(E)$  can be represented as product  $\gamma = \gamma_1^{-1} \cdot \gamma_2$  where  $\gamma_1, \gamma_2, s(\gamma_1) = s(\gamma_2) = 0$ ; define a map  $h'_E : \Omega(E) \rightarrow \Omega'(E)$  by setting  $h'_E(\gamma) = h(\gamma_1)^{-1} \cdot h(\gamma_2)$ . If  $\gamma_1^{-1}\gamma_2 = \gamma_1'^{-1}\gamma_2'$  then  $\gamma_1^{-1}\gamma_1' = \gamma_2\gamma_2'^{-1} \in \Omega_{0,0}(E)$ , and so

$$\begin{aligned} h(\gamma_1)^{-1} \cdot h(\gamma_2) &= h((\gamma_1\gamma_1'^{-1})\gamma_1')^{-1} \cdot h((\gamma_2\gamma_2'^{-1})\gamma_2') = \\ &= h((\gamma_1^{-1}\gamma_1') + \gamma_1')^{-1} \cdot h((\gamma_2\gamma_2'^{-1}) + \gamma_2') = \\ &= (h(\gamma_1^{-1}\gamma_1') + h(\gamma_1'))^{-1} \cdot h(\gamma_2\gamma_2'^{-1}) + h(\gamma_2') = \\ &= (h(\gamma_1'))^{-1} \cdot h(\gamma_2') = h(\gamma_1'^{-1}) \cdot h(\gamma_2'), \end{aligned}$$

which proves that  $h'$  is well-defined. Similar calculations check that  $h'_E$  preserves concatenation  $\cdot_E$ :  $h'(\gamma \cdot \gamma') = h'(\gamma) \cdot h'(\gamma')$  for arbitrary  $\gamma, \gamma' \in \Omega(E)$ .

We define  $h'_{E^n} = h'_{E^n} : \Omega(E^n) \rightarrow \Omega'(E^n)$  from an arbitrary Cartesian power  $\Omega(E^n) = \Omega(E)^n$  by  $h'_{E \times E}(\gamma \times \gamma') = h'_E(\gamma) \times h'_E(\gamma')$  etc.

4.1.6. *Checking that  $h'$  is a natural transformation of  $\underline{\Omega}$  into  $\underline{\Omega}' \circ \sigma$ .* To check that  $h'$  is a natural transformation of  $\underline{\Omega}$  to  $\underline{\Omega}' \circ \sigma$ , we need to check commutativity of the following diagrams (note that  $\sigma$  on the left-hand side is *not* a morphism but a functor!):

$$(2) \quad \begin{array}{ccc} E^n & \xrightarrow{\text{Pt}} & E^n(\overline{\mathbb{Q}}) \\ \sigma \downarrow & & \downarrow \sigma_{E^n} \\ E^n & \xrightarrow{\text{Pt}} & E^n(\overline{\mathbb{Q}}) \end{array} \quad \begin{array}{ccc} A(\overline{\mathbb{Q}}) & \xrightarrow{\text{Pt}(f)} & B(\overline{\mathbb{Q}}) \\ \sigma_A \downarrow & & \downarrow \sigma_B \\ A(\overline{\mathbb{Q}}) & \xrightarrow{\text{Pt}(f \circ \sigma)} & B(\overline{\mathbb{Q}}) \end{array}$$

$$(3) \quad \begin{array}{ccc} E^n & \xrightarrow{\Omega} & \Omega(E^n) \\ \sigma \downarrow & & \downarrow h'_{E^n} \\ E^n & \xrightarrow{\Omega'} & \Omega'(E^n) \end{array} \quad \begin{array}{ccc} \Omega(A) & \xrightarrow{\Omega(f)} & \Omega(B) \\ h'_A \downarrow & & \downarrow h'_B \\ \Omega'(A) & \xrightarrow{\Omega'(f \circ \sigma)} & \Omega'(B) \end{array}$$

$$(4) \quad \begin{array}{ccc} \Omega(A) & \xrightarrow{s_A} & \text{Pt}(A) \\ h'_A \downarrow & & \downarrow \sigma_A \\ \Omega'(A) & \xrightarrow{s'_A} & \text{Pt}(A) \end{array} \quad \begin{array}{ccc} \Omega(A) & \xrightarrow{t_A} & \text{Pt}(A) \\ h'_A \downarrow & & \downarrow \sigma_A \\ \Omega'(A) & \xrightarrow{t'_A} & \text{Pt}(A) \end{array}$$

$$(5) \quad \begin{array}{ccc} \Omega(A) \times \Omega(A) & \xrightarrow[\text{partial}]{\cdot_A} & \Omega(A) \\ h'_A \times h'_A \downarrow & & h'_A \downarrow \\ \Omega'(A) \times \Omega'(A) & \xrightarrow[\text{partial}]{\cdot'_A} & \Omega'(A) \end{array}$$

The first pair (2) expresses that  $h_A^{\text{Pt}} = (\sigma_A)_A$  is a natural transformation of set-valued functors from  $\text{Pt}$  to  $\text{Pt} \circ \sigma$ ; the diagrams are commutative just by definition of the action of  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  on category  $\mathfrak{E}$ . Analogously the second pair (3) expresses that  $h'$  is a natural transformation of set-valued functors  $\Omega$  to  $\Omega' \circ \sigma$ . The first diagram in (3) says that  $h'_{E^n}$  is a map from  $\Omega(E^n)$  to  $\Omega'(\sigma(E^n)) = \Omega'(\sigma(E^n))$ ; the second one in (3) expresses the linearity of  $h'$  with respect to the morphisms of  $E^n \rightarrow E^m$ ; that follows from  $\text{End}E$ -linearity of  $h : V \rightarrow V$ .

The third pair (4) expresses compatibility of the end-point functions and  $h'_A$ ; for  $A = E$ , this follows from the main property  $\sigma \circ s = s \circ h$  when restricted to  $W^n \subset \Omega(E^n)$ , and that  $h'$  preserves concatenation  $\cdot_A$ ; preservation of direct products allows us to extend this to arbitrary Cartesian power  $E^n$ . The diagram (5) expresses the fact that  $h'$  preserves concatenation; this is by the definition of  $h'$ .

This concludes the proof that  $h'$  is a natural transformation and that of derivation of Proposition 2 from Proposition 1.  $\square$

## 5. SHIMURA CURVES

Arithmetics of Shimura curves is well-studied. In particular, for Shimura curves, there is a quite explicit description of Galois action analogous to the results on the Galois action on the Tate module of an elliptic curve: for curves without complex multiplication it is a result of Ohta ([Oht74], also [Cla03, Theorem 123, p. 90]); for curves with complex multiplication this is implied by the explicit description of Galois group provided by the theory of complex multiplication. We thank A.Yafaev for pointing and explaining us those results. These results motivate us to make the following conjecture.

Let  $S$  be a connected Shimura curve defined over a number field  $k \subset \overline{\mathbb{Q}}$ , and let  $\mathfrak{SC}$  be full subcategory of  $\text{Var}/\overline{\mathbb{Q}}$  consisting of Cartesian powers of finite étale covers of  $S$ . Assume that  $S$  has a  $k$ -rational point  $O$ .

Recall we denote  $\text{Pt}_{\pi_1^{\text{top}}|\overline{\mathbb{Q}}}(V) = V(\overline{\mathbb{Q}})$  and  $\Omega_{\pi_1^{\text{top}}|\overline{\mathbb{Q}}}(V) = \{\gamma \in \pi_1^{\text{top}}(V(\mathbb{C})) : s(\gamma), t(\gamma) \in V(\overline{\mathbb{Q}})\}$  is the restriction of  $\pi_1^{\text{top}}$  to  $\overline{\mathbb{Q}}$ -rational points.

**Conjecture 10** (Universality of fundamental groupoid functor). *Let  $\underline{\Omega} = (\text{Pt}, \Omega, s_V, t_V, \cdot_V)$  be a functor from category  $\mathfrak{E}$  to  $\mathfrak{Groupoids}$  such that*

- (1) *the functor of points of is the functor of  $\overline{\mathbb{Q}}$ -rational points:*  
 $\text{Pt}(X) = X(\overline{\mathbb{Q}}) = \mathcal{M}or_{\mathfrak{E}}(O, X)$ ,  $X \in \mathfrak{E}$ .
- (2)  *$\underline{\Omega}$  preserves direct product:  $\underline{\Omega}(X \times Y) = \underline{\Omega}(X) \times \underline{\Omega}(Y)$ .*
- (3)  *$\underline{\Omega}$  has the unique path-lifting property along étale morphisms: for an étale morphism  $f : X \rightarrow Y$ , a path  $\gamma \in \Omega(Y)$  and a point  $x \in \text{Pt}(X)$  such that  $\text{Pt}(f)(x) = s(\gamma)$ , there exists a unique path  $\tilde{\gamma} \in \Omega(X)$  such that  $\Omega(f)(\tilde{\gamma}) = \gamma$  and  $s(\tilde{\gamma}) = x$ .*

*Assume further that*

- (4)  *$\underline{\Omega}(E)$  is a connected groupoid*
- (5) *there is an isomorphism*

$$\Omega_{0,0}(S) := \{\gamma \in \Omega(S) : s(\gamma) = t(\gamma) = 0\} \cong \pi_1^{\text{top}}(S(\mathbb{C}), O).$$

*Then there exists an automorphism  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that the functors  $\underline{\Omega}$  and  $\pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma : \mathfrak{E} \rightarrow \mathfrak{Groupoids}$  are naturally equivalent:*

$$\underline{\Omega} \cong \pi_1^{\text{top}}|_{\overline{\mathbb{Q}}} \circ \sigma$$

It is possible that only a weaker conclusion holds: there exists *finitely many* functors  $\mathbb{F}_1, \dots, \mathbb{F}_n$  satisfying (1)-(5) such that for any functor  $\underline{\Omega}$  satisfying (1)-(5) there exists  $\sigma \in \text{Gal}(\overline{\mathbb{Q}}/k)$  such that  $\underline{\Omega} \circ \sigma$  is naturally equivalent to one of  $\mathbb{F}_1, \dots, \mathbb{F}_n$ .

## REFERENCES

- [Bas72] M. Bashmakov. The cohomology of an abelian variety over a number field. *Russian Mathematical Surveys*, 27:25–70, 1972.
- [Ber88] Daniel Bertrand. Galois representations and transcendental numbers. In *New advances in transcendence theory (Durham, 1986)*, pages 37–55. Cambridge Univ. Press, Cambridge, 1988.
- [Cla03] Pete L. Clark. *Local and Global Moduli Spaces of Potentially Quaternionic Abelian Surfaces*. PhD thesis, Harvard University, April 2003.
- [Gav05] Misha Gavrilovich. *Model Theory of the Universal Covering Spaces of Complex Algebraic Varieties*. PhD thesis, Oxford University, 2005. submitted, available at <http://misha.uploads.net.ru/misha-thesis.pdf>.
- [Gro] Alexander Grothendieck. Pursuing stacks, *also known as* Long Letter to Quillen. <http://www.math.jussieu.fr/~leila/mathtexts.php>.
- [JR87] Olivier Jacquinet and Kenneth A. Ribet. Deficient points on extensions of abelian varieties by  $\mathbf{G}_m$ . *J. Number Theory*, 25(2):133–151, 1987.
- [Lan78] Serge Lang. *Elliptic curves: Diophantine analysis*, volume 231 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1978.
- [Lan83] Serge Lang. *Complex multiplication*. Springer, 1983.

- [Mum70] David Mumford. *Abelian varieties*. Tata Institute of Fundamental Research Studies in Mathematics, No. 5. Published for the Tata Institute of Fundamental Research, Bombay, 1970.
- [Oht74] M. Ohta. On  $l$ -adic representations of Galois groups obtain from certain two-dimensional abelian varieties. *J. Fac. Soc. Univ. Tokyo Sect. 1A Math.*, 21:299–308, 1974.
- [Rib79] Karl Ribet. Kummer theory on extensions of varieties by tori. *Duke Mathematical Journal*, 46, 1979.
- [Rib87] Kenneth A. Ribet. Cohomological realization of a family of 1-motives. *J. Number Theory*, 25(2):152–161, 1987.
- [Shi71] G. Shimura. *Introduction to the arithmetic theory of automorphic functions*. Princeton: Princeton University Press, 1971.
- [VK91] Vladimir Voevodsky and Mikhail Kapranov.  $\infty$ -groupoids and homotopy types. *Cah. Top. Géom. Diff. Cat.*, 32:29–46, 1991.
- [Zila] Boris Zilber. Covers of the multiplicative group of an algebraically closed field. <http://www.maths.ox.ac.uk/~zilber>.
- [Zilb] Boris Zilber. Logically perfect structures. slides, available at <http://www.maths.ox.ac.uk/~zilber>.

URL: [HTTP://MISHA.UPLOADS.NET.RU](http://MISHA.UPLOADS.NET.RU)